


Sexually Transmitted Infections Meded101



Across

- Antibiotic recommended by the CDC for the treatment of gonorrhea
- Antibiotic treatment for bacterial vaginosis that may be given orally as a tablet or intravaginally as a gel
- A firm, round, painless sore found at the site of infection during early primary syphilis
- Typically asymptomatic, but patients may experience burning, painful urination, watery discharge, and/or abdominal pain with this STI

Down

- This form of penicillin G should be used to treat early primary syphilis
- Using this during sexual activity can help prevent the spread of STIs
- Reinfection with trichomoniasis is common, so patients should consider re-testing in _____ months
- Antibiotic recommended by the CDC for the treatment of chlamydia
- HPV vaccine covering two types of HIV and only indicated for females

11. Patients may present with painful urination, thick discharge, abdominal pain, and/or testicular swelling with this STI
13. Bacterial vaginosis results from an overgrowth of _____ bacteria that produce volatile amines (fish-smelling discharge)
14. Common adverse effect of penicillin antibiotics
18. Preferred antibiotic for the treatment of chlamydia in pregnancy
20. Administering penicillin G benzathine or procaine by this route can be fatal
21. Ceftriaxone for gonorrhea should be administered by this route
22. When acyclovir is used to treat the first episode of herpes, it should be dosed as 400 mg by mouth _____ times daily
23. Treatment for pelvic inflammatory disease (PID) may be given intravenously, intramuscularly, or _____
24. Herpes may be spread through _____, genital secretions, or direct contact with skin in the oral or genital areas
9. Class of anti-infectives used to treat HIV
10. Most prevalent non-viral STI caused by *Trichomonas vaginalis*
12. Class of anti-infectives used to treat herpes
15. All partners within the past _____ days should be treated for syphilis
16. Twenty-eight _____ is the appropriate duration of treatment for HIV post-exposure prophylaxis
17. All partners within the past _____ days should be treated for chlamydia
19. Virus responsible for causing genital warts and cervical, anal, and penile cancers